

# QCDF Amplitudes from SU(3) Symmetries

Gilberto Tetlalmatzi-Xolocotzi


*Based on: T. Huber and GTX, 2111.06418  
Eur.Phys.J.C 81 (2021) 7, 658*

**CPPS, Theoretische Physik 1,  
Universität Siegen**



# Non-leptonic B meson decays

We are interested in *B meson decays into pairs of light pseudoscalar mesons*



$$B \rightarrow PP$$

The light pseudoscalar mesons are bound states of light quarks  $[u, d, s]$  (SU(3) symmetry)

$$B = (B^+, B_d^0, B_s^0)$$


$$q_i \otimes \bar{q}_j \rightarrow 3 \otimes \bar{3} = 8 \oplus 1$$

$$i, j \in [u, d, s]$$

$$P \rightarrow M = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} & \frac{\eta_q}{\sqrt{2}} + \frac{\eta'_q}{\sqrt{2}} & \pi^- & K^- \\ \pi^+ & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_q}{\sqrt{2}} + \frac{\eta'_q}{\sqrt{2}} & \bar{K}^0 & \\ K^+ & K^0 & \eta_s + \eta'_s & \end{pmatrix}$$

# Non-leptonic B meson decays

We are interested in *B meson* decays into pairs of *light pseudoscalar mesons*



$$b \rightarrow u \bar{u} q$$

$$q = d, s$$

Several possible decay channels

$$\overline{B}^- \rightarrow \pi^0 \pi^-$$

$$\overline{B}^- \rightarrow \pi^- \eta_8$$

$$\overline{B}^- \rightarrow \pi^- \eta_1$$

$$\overline{B}^- \rightarrow K^0 K^-$$

$$\overline{B}^0 \rightarrow \pi^+ \pi^-$$

$$\overline{B}^0 \rightarrow \pi^0 \pi^0$$

$$\overline{B}^0 \rightarrow \pi^0 \eta_8$$

$$\overline{B}^0 \rightarrow \pi^0 \eta_1$$

$$\overline{B}^0 \rightarrow K^+ K^-$$

$$\overline{B}^0 \rightarrow K^0 \overline{K}^0$$

$$\overline{B}^0 \rightarrow \eta_8 \eta_8$$

$$\overline{B}^0 \rightarrow \eta_8 \eta_1$$

$$\overline{B}^0 \rightarrow \eta_1 \eta_1$$

$$\overline{B}_s^0 \rightarrow \pi^0 K^0$$

$$\overline{B}_s^0 \rightarrow \pi^- K^+$$

$$\overline{B}_s^0 \rightarrow K^0 \eta_8$$

$$\overline{B}_s^0 \rightarrow K^0 \eta_1$$

$$B^- \rightarrow \pi^0 K^-$$

$$B^- \rightarrow \pi^- \overline{K}^0$$

$$B^- \rightarrow K^- \eta_8$$

$$B^- \rightarrow K^- \eta_1$$

$$\overline{B}^0 \rightarrow \pi^+ K^-$$

$$\overline{B}^0 \rightarrow \pi^0 \overline{K}^0$$

$$\overline{B}^0 \rightarrow \overline{K}^0 \eta_8$$

$$\overline{B}^0 \rightarrow \overline{K}^0 \eta_1$$

$$\overline{B}_s^0 \rightarrow \pi^+ \pi^-$$

$$\overline{B}_s^0 \rightarrow \pi^0 \pi^0$$

$$\overline{B}_s^0 \rightarrow \pi^0 \eta_1$$

$$\overline{B}_s^0 \rightarrow K^+ K^-$$

$$\overline{B}_s^0 \rightarrow K^0 \overline{K}^0$$

$$\overline{B}_s^0 \rightarrow \eta_8 \eta_8$$

$$\overline{B}_s^0 \rightarrow \eta_8 \eta_1$$

$$\overline{B}_s^0 \rightarrow \eta_1 \eta_1$$

# Topological decomposition

Consider the process  $B \rightarrow PP$

where  $P$  is a charmless pseudoscalar meson

*The physical amplitude can be decomposed as*

$$\mathcal{A}^{TDA} = i \frac{G_F}{\sqrt{2}} \left[ \mathcal{T}^{TDA} + \mathcal{P}^{TDA} \right]$$

$$\lambda_p^{(q)} = V_{pb} V_{pq}^* \quad \lambda_u^{(q)} \quad \lambda_t^{(q)} \quad q = d, s$$

$$\lambda_u^{(q)} + \lambda_c^{(q)} + \lambda_t^{(q)} = 0$$

# Topological decomposition

Consider the process  $B \rightarrow PP$

where  $P$  is a charmless pseudoscalar meson

*The physical amplitude can be decomposed as*

$$\mathcal{A}^{TDA} = i \frac{G_F}{\sqrt{2}} \left[ \mathcal{T}^{TDA} + \mathcal{P}^{TDA} \right]$$

$$\lambda_p^{(q)} = V_{pb} V_{pq}^* \quad \lambda_u^{(q)} \quad \lambda_t^{(q)} \quad q = d, s$$

$$\lambda_u^{(q)} + \lambda_c^{(q)} + \lambda_t^{(q)} = 0$$

# Topological decomposition

$$\begin{aligned}
 \mathcal{T}^{TDA} = & \underline{T} B_i(M)_j^i \bar{H}_k^{jl}(M)_l^k + \underline{C} B_i(M)_j^i \bar{H}_k^{lj}(M)_l^k + \underline{A} B_i \bar{H}_j^{il}(M)_k^j (M)_l^k \\
 & + \underline{E} B_i \bar{H}_j^{li}(M)_k^j (M)_l^k + \underline{T_{ES}} B_i \bar{H}_l^{ij}(M)_j^l (M)_k^k + \underline{T_{AS}} B_i \bar{H}_l^{ji}(M)_j^l (M)_k^k \\
 & + \underline{T_S} B_i(M)_j^i \bar{H}_l^{lj}(M)_k^k + \underline{T_{PA}} B_i \bar{H}_l^{li}(M)_k^j (M)_j^k + \underline{T_P} B_i(M)_j^i (M)_k^j \bar{H}_l^{lk} \\
 & + \underline{T_{SS}} B_i \bar{H}_l^{li}(M)_j^j (M)_k^k,
 \end{aligned}$$

SU(3) Flavour

$[u, d, s]$


$$B = (B^+, B_d^0, B_s^0)$$

$$M = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_q}{\sqrt{2}} + \frac{\eta'_q}{\sqrt{2}} & \pi^- & K^- \\ \pi^+ & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_q}{\sqrt{2}} + \frac{\eta'_q}{\sqrt{2}} & \bar{K}^0 \\ K^+ & K^0 & \eta_s + \eta'_s \end{pmatrix}$$

$$\bar{H}_1^{12} = \lambda_u^{(d)}, \quad \bar{H}_1^{13} = \lambda_u^{(s)},$$

# Topological decomposition

$$\begin{aligned}
 \mathcal{T}^{TDA} = & \underline{T} B_i(M)_j^i \bar{H}_k^{jl}(M)_l^k + \underline{C} B_i(M)_j^i \bar{H}_k^{lj}(M)_l^k + \underline{A} B_i \bar{H}_j^{il}(M)_k^j (M)_l^k \\
 & + \underline{E} B_i \bar{H}_j^{li}(M)_k^j (M)_l^k + \underline{T_{ES}} B_i \bar{H}_l^{ij}(M)_j^l (M)_k^k + \underline{T_{AS}} B_i \bar{H}_l^{ji}(M)_j^l (M)_k^k \\
 & + \underline{T_S} B_i(M)_j^i \bar{H}_l^{lj}(M)_k^k + \underline{T_{PA}} B_i \bar{H}_l^{li}(M)_k^j (M)_j^k + \underline{T_P} B_i(M)_j^i (M)_k^j \bar{H}_l^{lk} \\
 & + \underline{T_{SS}} B_i \bar{H}_l^{li}(M)_j^j (M)_k^k,
 \end{aligned}$$



# SU(3)-Irreducible decomposition

$$\begin{aligned}
 \mathcal{T}^{IRA} = & \underline{A_3^T} B_i (\bar{H}_3)^i (M)_k^j (M)_j^k + \underline{C_3^T} B_i (M)_j^i (M)_k^j (\bar{H}_3)^k + \underline{B_3^T} B_i (\bar{H}_3)^i (M)_k^k (M)_j^j \\
 & + \underline{D_3^T} B_i (M)_j^i (\bar{H}_3)^j (M)_k^k + \underline{A_6^T} B_i (H_6)_k^{ij} (M)_j^l (M)_l^k + \underline{C_6^T} B_i (M)_j^i (\bar{H}_6)_k^{jl} (M)_l^k \\
 & + \underline{B_6^T} B_i (\bar{H}_6)_k^{ij} (M)_j^k (M)_l^l + \underline{A_{15}^T} B_i (\bar{H}_{15})_k^{ij} (M)_j^l (M)_l^k + \underline{C_{15}^T} B_i (M)_j^i (\bar{H}_{15})_l^{jk} (M)_k^l \\
 & + \underline{B_{15}^T} B_i (\bar{H}_{15})_k^{ij} (M)_j^k (M)_l^l.
 \end{aligned}$$

## SU(3) irreducible decomposition

$$\bar{H}_k^{ij} = \frac{1}{8} (H_{15})_k^{ij} + \frac{1}{4} (H_6)_k^{ij} - \frac{1}{8} (H_3)^i \delta_k^j + \frac{3}{8} (H_{3'})^j \delta_k^i$$



# SU(3)-Irreducible decomposition

$$\begin{aligned}
 \mathcal{T}^{IRA} = & \underline{A_3^T} B_i (\bar{H}_3)^i (M)_k^j (M)_j^k + \underline{C_3^T} B_i (M)_j^i (M)_k^j (\bar{H}_3)^k + \underline{B_3^T} B_i (\bar{H}_3)^i (M)_k^j (M)_j^k \\
 & + \underline{D_3^T} B_i (M)_j^i (\bar{H}_3)^j (M)_k^k + \underline{A_6^T} B_i (H_6)^{ij} (M)_j^l (M)_l^k + \underline{C_6^T} B_i (M)_j^i (\bar{H}_6)^{jl} (M)_l^k \\
 & + \underline{B_6^T} B_i (\bar{H}_6)^{ij} (M)_j^k (M)_l^l + \underline{A_{15}^T} B_i (\bar{H}_{15})^{ij} (M)_j^l (M)_l^k + \underline{C_{15}^T} B_i (M)_j^i (\bar{H}_{15})^{jk} (M)_k^l \\
 & + \underline{B_{15}^T} B_i (\bar{H}_{15})^{ij} (M)_j^k (M)_l^l.
 \end{aligned}$$

## Topological to SU(3)

X.-G. He and W. Wang: 1803.04227

$$A_3^T = -\frac{A}{8} + \frac{3E}{8} + T_{PA},$$

$$B_3^T = T_{SS} + \frac{3T_{AS} - T_{ES}}{8},$$

$$C_3^T = \frac{1}{8}(3A - C - E + 3T) + T_P,$$

$$D_3^T = T_S + \frac{1}{8}(3C - T_{AS} + 3T_{ES} - T)$$

$$A_6^T = \frac{1}{4}(A - E),$$

$$B_6^T = \frac{1}{4}(T_{ES} - T_{AS}),$$

$$C_6^T = \frac{1}{4}(-C + T),$$

$$A_{15}^T = \frac{A + E}{8},$$

$$B_{15}^T = \frac{T_{ES} + T_{AS}}{8},$$

$$C_{15}^T = \frac{C + T}{8},$$

# SU(3) amplitudes from data

The physical amplitudes can be expressed as linear combinations of the SU(3) sub-amplitudes

Channel	$A_3^T$	$C_3^T$	$A_6^T$	$C_6^T$	$A_{15}^T$	$C_{15}^T$	$B_3^T$	$B_6^T$	$B_{15}^T$	$D_3^T$
$B^- \rightarrow \pi^0 \pi^-$	0	0	0	0	0	$4\sqrt{2}$	0	0	0	0
$B^- \rightarrow K^0 K^-$	0	1	1	-1	3	-1	0	0	0	0
$B^0 \rightarrow \pi^+ \pi^-$	2	1	-1	1	1	3	0	0	0	0
$B^0 \rightarrow \pi^0 \pi^0$	2	1	-1	1	1	-5	0	0	0	0
$B^0 \rightarrow K^+ K^-$	2	0	0	0	2	0	0	0	0	0
$B^0 \rightarrow K^0 \bar{K}^0$	2	1	1	-1	-3	-1	0	0	0	0
$B_s \rightarrow \pi^0 K^0$	0	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$\frac{5}{\sqrt{2}}$	0	0	0	0
$B_s \rightarrow \pi^- K^+$	0	1	-1	1	-1	3	0	0	0	0
$B^- \rightarrow \pi^0 K^-$	0	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{3}{\sqrt{2}}$	$\frac{7}{\sqrt{2}}$	0	0	0	0
$B^- \rightarrow \pi^- K^0$	0	1	1	-1	3	-1	0	0	0	0
$B^0 \rightarrow \pi^+ K^-$	0	1	-1	1	-1	3	0	0	0	0
$B^0 \rightarrow \pi^0 K^0$	0	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$\frac{5}{\sqrt{2}}$	0	0	0	0
$B_s \rightarrow \pi^+ \pi^-$	2	0	0	0	2	0	0	0	0	0
$B_s \rightarrow \pi^0 \pi^0$	2	0	0	0	2	0	0	0	0	0
$B_s \rightarrow K^+ K^-$	2	1	-1	1	1	3	0	0	0	0
$B_s \rightarrow K^0 \bar{K}^0$	2	1	1	-1	-3	-1	0	0	0	0

# SU(3) amplitudes from data

Extract the SU(3) amplitudes by fitting to data

$$\Gamma(\bar{B} \rightarrow M_1 M_2) = \frac{S}{16\pi M_B} |\mathcal{A}_{B \rightarrow M_1 M_2}|^2$$

$$S=1 \quad \text{if} \quad M_1 \neq M_2 \quad \quad S=1/2 \quad \text{if} \quad M_1 = M_2$$

Observables:

Branching fractions  $\mathcal{B}(\bar{B} \rightarrow \bar{f}) = \frac{1}{2} \tau_B [\Gamma(\bar{B} \rightarrow \bar{f}) + \Gamma(B \rightarrow f)]$

CP Asymmetries  $\mathcal{A}_{\text{CP}}(\bar{B} \rightarrow \bar{f}) = \frac{\Gamma(\bar{B} \rightarrow \bar{f}) - \Gamma(B \rightarrow f)}{\Gamma(\bar{B} \rightarrow \bar{f}) + \Gamma(B \rightarrow f)}$

# SU(3) amplitudes from data

Perform a  $\chi^2$  fit  $\chi^2 = \sum_i \left( \frac{O_i^{\text{Theo}} - O_i^{\text{Exp}}}{\sigma_i^{\text{Exp}}} \right)^2$

10 Tree complex amplitudes

$$A_3^T, C_3^T, A_6^T, C_6^T, A_{15}^T, C_{15}^T, B_3^T, B_6^T, B_{15}^T, D_3^T$$

and 10 Penguin complex amplitudes (replace T for P above)

The combinations  $C_6^T - \underline{A_6^T}$  and  $B_6^T + \underline{A_6^T}$  always appear together (analogously for penguins)

Redefine

$$\begin{aligned} C_6^T - A_6^T &\rightarrow C_6^T & C_6^P - A_6^P &\rightarrow C_6^P \\ B_6^T + A_6^T &\rightarrow B_6^T & B_6^P + A_6^P &\rightarrow B_6^P \end{aligned}$$

Absorb a global phase by taking  $C_3^P$  as a real parameter

35 parameters +  $\theta_{FKS}$  = 36 parameters to fit.

# SU(3) amplitudes from data

Best fit point (modulus in  $\text{GeV}^3$ )

<u><math> A_3^T  = 0.029,</math></u>	$\delta_{A_3^T} = -3.083,$	$ C_3^T  = 0.258,$	$\delta_{C_3^T} = -0.105,$
$ C_6^T  = 0.235,$	$\delta_{C_6^T} = -0.079,$	<u><math> A_{15}^T  = 0.029,</math></u>	$\delta_{A_{15}^T} = -3.083,$
$ C_{15}^T  = 0.151,$	$\delta_{C_{15}^T} = 0.061,$	<u><math> B_3^T  = 0.034,</math></u>	$\delta_{B_3^T} = 3.087$
<u><math> B_6^T  = 0.033,</math></u>	$\delta_{B_6^T} = -0.286,$	<u><math> B_{15}^T  = 0.008,</math></u>	$\delta_{B_{15}^T} = -1.892$
$ D_3^T  = 0.055,$	$\delta_{D_3^T} = 2.942,$		
<u><math> A_3^P  = 0.014,</math></u>	$\delta_{A_3^P} = -1.328,$	$ C_6^P  = 0.145,$	$\delta_{C_6^P} = -2.881,$
<u><math> A_{15}^P  = 0.003,</math></u>	$\delta_{A_{15}^P} = 2.234,$	<u><math> C_{15}^P  = 0.003,</math></u>	$\delta_{C_{15}^P} = -0.608,$
<u><math> B_3^P  = 0.043,</math></u>	$\delta_{B_3^P} = 2.367,$	<u><math> B_6^P  = 0.099,</math></u>	$\delta_{B_6^P} = 0.353,$
<u><math> B_{15}^P  = 0.031,</math></u>	$\delta_{B_{15}^P} = -0.690,$	$ D_3^P  = 0.030,$	$\delta_{D_3^P} = 0.477,$
$ C_3^P  = 0.008,$	$\theta_{FKS} = 0.628.$		

Annihilation amplitudes below 10%.

$$\chi^2/d.o.f. = 0.851$$

# Fit-Results: Branching fractions

Channel	Branching ratio in units of $10^{-6}$		Channel	Branching ratio in units of $10^{-6}$	
	Experimental	Theoretical		Experimental	Theoretical
$B^- \rightarrow \pi^0 \pi^-$	$5.5 \pm 0.4$	$6.04^{+2.42}_{-2.51}$	$B^- \rightarrow \eta \pi^-$	$4.02 \pm 0.27$	$3.80^{+1.25}_{-1.55}$
$B^- \rightarrow K^0 K^-$	$1.31 \pm 0.17$	$1.36^{+0.17}_{-0.16}$	$B^- \rightarrow \eta' \pi^-$	$2.7 \pm 0.9$	$3.55^{+4.49}_{-1.67}$
$\bar{B}^0 \rightarrow \pi^+ \pi^-$	$5.12 \pm 0.19$	$6.31^{+0.61}_{-0.50}$	$\bar{B}^0 \rightarrow \eta \pi^0$	$0.41 \pm 0.17$	$0.41^{+8.90}_{-4.08}$
$\bar{B}^0 \rightarrow \pi^0 \pi^0$	$1.59 \pm 0.26$	$1.01^{+1.30}_{-0.51}$	$\bar{B}^0 \rightarrow \eta' \pi^0$	$1.2 \pm 0.6$	$1.20^{+3.62}_{-1.19}$
$\bar{B}^0 \rightarrow K^+ K^-$	$0.078 \pm 0.015$	$0.13^{+0.08}_{-0.07}$	$\bar{B}_s \rightarrow \eta K^0$	Not available	$0.13^{+0.11}_{-0.08}$
$\bar{B}^0 \rightarrow K^0 \bar{K}^0$	$1.21 \pm 0.16$	$1.13^{+0.83}_{-0.91}$	$\bar{B}_s \rightarrow \eta' K^0$	Not available	$6.65^{+1.48}_{-1.65}$
$\bar{B}_s \rightarrow \pi^- K^+$	$5.8 \pm 0.7$	$7.75^{+0.63}_{-0.09}$	$B^- \rightarrow \eta K^-$	$2.4 \pm 0.4$	$2.34^{+1.39}_{-1.67}$
$B^- \rightarrow \pi^0 K^-$	$12.9 \pm 0.5$	$12.78^{+1.75}_{-1.94}$	$B^- \rightarrow \eta' K^-$	$70.4 \pm 2.5$	$70.82^{+11.16}_{-11.53}$
$B^- \rightarrow \pi^- \bar{K}^0$	$23.7 \pm 0.8$	$23.85^{+2.23}_{-2.31}$	$\bar{B}^0 \rightarrow \eta K^0$	$1.23 \pm 0.27$	$1.38^{+1.15}_{-0.36}$
$\bar{B}^0 \rightarrow \pi^+ K^-$	$19.6 \pm 0.5$	$19.47^{+1.72}_{-2.24}$	$\bar{B}^0 \rightarrow \eta' K^0$	$6.6 \pm 0.4$	$6.65^{+1.48}_{-1.65}$
$\bar{B}^0 \rightarrow \pi^0 \bar{K}^0$	$9.9 \pm 0.5$	$10.17^{+2.00}_{-2.30}$	$\bar{B}_s \rightarrow \eta \pi^0$	$< 10^3$	$31.15^{+39.05}_{-31.14}$
$\bar{B}_s \rightarrow \pi^+ \pi^-$	$0.7 \pm 0.1$	$0.57^{+0.40}_{-0.42}$	$\bar{B}_s \rightarrow \eta' \pi^0$	Not available	$11.13^{+74.75}_{-11.12}$
$\bar{B}_s \rightarrow \pi^0 \pi^0$	$< 210$	$0.28^{+0.20}_{-0.21}$	$\bar{B}^0 \rightarrow \eta \eta$	$< 1$	$0.30^{+0.70}_{-0.30}$
$\bar{B}_s \rightarrow K^+ K^-$	$26.6 \pm 2.2$	$20.63^{+6.80}_{-8.09}$	$\bar{B}_s \rightarrow \eta \eta$	$< 1.5 \times 10^3$	$2.58^{+36.53}_{-2.57}$
$\bar{B}_s \rightarrow K^0 \bar{K}^0$	$20 \pm 6$	$24.64^{+18.84}_{-21.14}$	$\bar{B}^0 \rightarrow \eta' \eta'$	$< 1.7$	$1.14^{+0.57}_{-1.07}$
$\bar{B}_s \rightarrow \pi^0 K^0$	Not available	$0.71^{+1.47}_{-0.27}$	$\bar{B}_s \rightarrow \eta' \eta'$	$33 \pm 7$	$33.00^{+24.52}_{-31.74}$
			$\bar{B}^0 \rightarrow \eta' \eta$	$< 1.2$	$0.61^{+0.59}_{-0.60}$
			$\bar{B}_s \rightarrow \eta' \eta$	Not available	$0.61^{+0.59}_{-0.60}$





Experimental results from PDG Live

# Fit-Results: CP Asymmetries

Channel	CP asymmetries in percent		Channel	CP asymmetries in percent	
	Experimental	Theoretical		Experimental	Theoretical
$B^- \rightarrow \pi^0 \pi^-$	$3 \pm 4$	$5.45^{+22.02}_{-20.60}$	$B^- \rightarrow \eta \pi^-$	$-14 \pm 7$	$-11.37^{+14.49}_{-26.90}$
$B^- \rightarrow K^0 K^-$	$4 \pm 14$	$18.82^{+36.93}_{-30.83}$	$B^- \rightarrow \eta' \pi^-$	$6 \pm 16$	$4.71^{+59.79}_{-57.97}$
$\bar{B}^0 \rightarrow \pi^+ \pi^-$	$32 \pm 4$	$35.01^{+3.19}_{-22.29}$	$\bar{B}_s \rightarrow \eta K^0$	$< 0.1$	$0.10^{+0.00}_{-100.07}$
$\bar{B}^0 \rightarrow \pi^0 \pi^0$	$33 \pm 22$	$-10.58^{+40.69}_{-89.40}$	$\bar{B}_s \rightarrow \eta' K^0$	Not available	$-0.58^{+100.57}_{-79.58}$
$\bar{B}^0 \rightarrow K^0 \bar{K}^0$	$-60 \pm 70$	$-6.88^{+85.39}_{-81.37}$	$B^- \rightarrow \eta K^-$	$-37 \pm 8$	$-42.23^{+42.23}_{-16.00}$
$\bar{B}_s \rightarrow \pi^- K^+$	$22.1 \pm 1.5$	$20.84^{+2.39}_{-2.57}$	$B^- \rightarrow \eta' K^-$	$0.4 \pm 1.1$	$0.63^{+3.98}_{-4.30}$
$B^- \rightarrow \pi^0 K^-$	$3.7 \pm 2.1$	$3.72^{+7.19}_{-4.35}$	$\bar{B}^0 \rightarrow \eta K^0$	Not available	$-0.01^{+40.07}_{-0.02}$
$B^- \rightarrow \pi^- K^0$	$-1.7 \pm 1.6$	$-1.08^{+1.76}_{-2.32}$	$\bar{B}^0 \rightarrow \eta' K^0$	$-6 \pm 4$	$0.03^{+4.82}_{-11.69}$
$\bar{B}^0 \rightarrow \pi^+ K^-$	$-8.3 \pm 0.4$	$-8.38^{+8.38}_{-1.01}$	$\bar{B}^0 \rightarrow \eta \pi^0$	Not available	$-27.39^{+127.11}_{-72.58}$
$\bar{B}^0 \rightarrow \pi^0 \bar{K}^0$	$0 \pm 13$	$-0.97^{+19.35}_{-3.20}$	$\bar{B}^0 \rightarrow \eta' \pi^0$	Not available	$-43.67^{+143.63}_{-56.33}$
$\bar{B}_s \rightarrow K^+ K^-$	$-14 \pm 11$	$-10.58^{+10.58}_{-3.60}$	$\bar{B}_s \rightarrow \eta \pi^0$	Not available	$0.88^{+94.98}_{-98.70}$
$\bar{B}_s \rightarrow \pi^+ \pi^-$	Not available	$17.56^{+11.84}_{-38.25}$	$\bar{B}_s \rightarrow \eta' \pi^0$	Not available	$1.57^{+77.56}_{-95.66}$
$\bar{B}_s \rightarrow \pi^0 \pi^0$	Not available	$17.56^{+11.84}_{-38.25}$	$\bar{B}^0 \rightarrow \eta \eta$	Not available	$3.46^{+96.50}_{-103.45}$
$\bar{B}_s \rightarrow K^0 \bar{K}^0$	Not available	$0.31^{+5.07}_{-4.59}$	$\bar{B}_s \rightarrow \eta \eta$	Not available	$14.29^{+76.81}_{-113.09}$
$\bar{B}^0 \rightarrow K^+ K^-$	Not available	$-78.45^{+161.99}_{-20.78}$	$\bar{B}^0 \rightarrow \eta' \eta'$	Not available	$42.41^{+57.55}_{-142.41}$
$\bar{B}_s \rightarrow \pi^0 K^0$	Not available	$13.74^{+29.49}_{-113.73}$	$\bar{B}_s \rightarrow \eta' \eta'$	Not available	$-2.05^{+15.29}_{-13.44}$
			$\bar{B}^0 \rightarrow \eta' \eta$	Not available	$-12.32^{+112.32}_{-87.67}$
			$\bar{B}_s \rightarrow \eta' \eta$	Not available	$3.43^{+96.36}_{-103.22}$

*Experimental results from PDG Live*

# SU(3) Confidence Regions






# QCF Factorization decomposition

*The topological and  $SU(3)$  invariant descriptions are just parametrizations of the decay amplitudes*

*A first principle technique to perform these calculations is QCD-Factorization*

*Beneke et al: 9905312*

*Beneke et al: 0308039*



*Naive Factorization*

$$\langle K \pi | Q | B \rangle \sim F_{B \rightarrow K} f_{\pi}$$


# QCF Factorization decomposition

*The topological and  $SU(3)$  invariant descriptions are just parametrizations of the decay amplitudes*

*A first principle technique to perform these calculations is QCD-Factorization*

*Beneke et al: 9905312*

*Beneke et al: 0308039*




*Naive Factorization*

$$\langle K \pi | Q | B \rangle \sim F_{B \rightarrow K} f_{\pi}$$

# QCF Factorization decomposition

*Naive factorization special case of*

$$\langle M_1 M_2 | \hat{Q}_i | B \rangle = \sum_j F_j^{B \rightarrow M_1}(0) \int_0^1 du T_{ij}^I(u) \Phi_{M_2}(u) + (M_1 \leftrightarrow M_2) \\ + \int_0^1 d\xi du dv T_i^{II}(\xi, u, v) \Phi_B(\xi) \Phi_{M_1}(v) \Phi_{M_2}(u).$$



*QCD-Factorization offers a systematic way to disentangle short from long distance physics considering  $\Lambda_{QCD} \ll m_b$*

# QCF Factorization decomposition

$$\begin{aligned}
 A^{\text{QCDF}} = & i \frac{G_F}{\sqrt{2}} \sum_{p=u,c} A_{M_1 M_2} \left\{ \underbrace{B M_1 \left( \alpha_1 \delta_{pu} \hat{U} + \alpha_4^p \hat{I} + \alpha_{4,EW}^p \hat{Q} \right)}_{\text{blue bar}} M_2 \Lambda_p \right. \\
 & + \underbrace{B M_1 \Lambda_p \cdot \text{Tr} \left[ \left( \alpha_2 \delta_{pu} \hat{U} + \alpha_3^p \hat{I} + \alpha_{3,EW}^p \hat{Q} \right) M_2 \right]}_{\text{blue bar}} \\
 & + B \left( \beta_2 \delta_{pu} \hat{U} + \beta_3^p \hat{I} + \beta_{3,EW}^p \hat{Q} \right) M_1 M_2 \Lambda_p \\
 & + \underbrace{B \Lambda_p \cdot \text{Tr} \left[ \left( \beta_1 \delta_{pu} \hat{U} + \beta_4^p \hat{I} + b_{4,EW}^p \hat{Q} \right) M_1 M_2 \right]}_{\text{blue bar}} \\
 & + B \left( \beta_{S2} \delta_{pu} \hat{U} + \beta_{S3}^p \hat{I} + \beta_{S3,EW}^p \hat{Q} \right) M_1 \Lambda_p \cdot \text{Tr} M_2 \\
 & \left. + \underbrace{B \Lambda_p \cdot \text{Tr} \left[ \left( \beta_{S1} \delta_{pu} \hat{U} + \beta_{S4}^p \hat{I} + b_{S4,EW}^p \hat{Q} \right) M_1 \right]}_{\text{blue bar}} \cdot \text{Tr} M_2 \right\}
 \end{aligned}$$

$$\Lambda_p = \begin{pmatrix} 0 \\ \lambda_p^{(d)} \\ \lambda_p^{(s)} \end{pmatrix},$$

$$\hat{U} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$


$$\hat{Q} = \frac{3}{2} Q = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix},$$

$$\hat{I} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$A_{M_1 M_2} = M_B^2 F_0^{B \rightarrow M_1}(0) f_{M_2}$$

# QCF Factorization decomposition


$$\begin{aligned}
 \mathcal{A}^{\text{QCDF}} = & i \frac{G_F}{\sqrt{2}} \sum_{p=u,c} A_{M_1 M_2} \left\{ \underbrace{B M_1 \left( \alpha_1 \delta_{pu} \hat{U} + \alpha_4^p \hat{I} + \alpha_{4,EW}^p \hat{Q} \right)}_{\text{Factorizable}} M_2 \Lambda_p \right. \\
 & + \underbrace{B M_1 \Lambda_p \cdot \text{Tr} \left[ \left( \alpha_2 \delta_{pu} \hat{U} + \alpha_3^p \hat{I} + \alpha_{3,EW}^p \hat{Q} \right) M_2 \right]}_{\text{Factorizable}} \\
 & + B \left( \beta_2 \delta_{pu} \hat{U} + \beta_3^p \hat{I} + \beta_{3,EW}^p \hat{Q} \right) M_1 M_2 \Lambda_p \\
 & + \underbrace{B \Lambda_p \cdot \text{Tr} \left[ \left( \beta_1 \delta_{pu} \hat{U} + \beta_4^p \hat{I} + b_{4,EW}^p \hat{Q} \right) M_1 M_2 \right]}_{\text{Factorizable}} \\
 & + B \left( \beta_{S2} \delta_{pu} \hat{U} + \beta_{S3}^p \hat{I} + \beta_{S3,EW}^p \hat{Q} \right) M_1 \Lambda_p \cdot \text{Tr} M_2 \\
 & \left. + \underbrace{B \Lambda_p \cdot \text{Tr} \left[ \left( \beta_{S1} \delta_{pu} \hat{U} + \beta_{S4}^p \hat{I} + b_{S4,EW}^p \hat{Q} \right) M_1 \right]}_{\text{Factorizable}} \cdot \text{Tr} M_2 \right\}
 \end{aligned}$$




→  $\beta$

Weak annihilation  
contributions  
are non-factorizable

# QCF Factorization decomposition



Weak annihilation contributions are non-factorizable



One of the main drawbacks of QCDF

These contributions are power suppressed

$$\Lambda_{QCD}/m_b$$

To address this problem educated Ansatz are made

$$X_A = \left(1 + \rho_A e^{i\phi_A}\right) \ln \frac{m_B}{\Lambda_h}$$

$$0 < \rho_A < 1$$

$$\Lambda_h \approx \mathcal{O}(\Lambda_{QCD})$$

*WHAT CAN WE LEARN ABOUT THE  
ANNIHILATION CONTRIBUTIONS FROM  
DATA?*

*CAN WE PROFIT FROM THE SU(3)  
INVARIANT FITS?*

*TO ACHIEVE THIS FIRST ESTABLISH A  
DICTIONARY BETWEEN SU(3) AND THE  
QCDF DECOMPOSITION OF THE  
PHYSICAL AMPLITUDES*

# QCF Factorization-Topological Equivalence

Equivalence between the QCF and the topological amplitudes

Decompose the matrix Q in terms of U and I  $\hat{Q} = \frac{3}{9}\hat{U} - \frac{1}{9}\hat{I}$

Use the  $\lambda_u^{(q)}$  and  $\lambda_t^{(q)}$  factors  $\Lambda_t = -\Lambda_u - \Lambda_c$

$$\mathcal{A}^{\text{QCDF}} = i\frac{G_F}{\sqrt{2}}A_{M_1M_2} \left\{ B_i M_j^i(\tilde{C}_1)_k^{jl} M_l^k + B_i M_j^i(\tilde{C}_2)_k^{lj} M_l^k + B_i(\tilde{C}_3)_k^{ij} M_l^k M_j^l \right. \\ \left. + B_i(\tilde{C}_4)_k^{li} M_r^k M_l^r + B_i(\tilde{C}_5)_k^{ij} M_i^k M_l^l + B_i(\tilde{C}_6)_k^{ji} M_j^k M_l^l \right\}$$

$$\tilde{C}_r = \left[ \tilde{T} + \frac{3}{2}\tilde{P}_2^u - \frac{3}{2}\tilde{P}_2^c \right] \hat{U} \otimes \Lambda_u + \left[ \tilde{P}_1^u - \tilde{P}_1^c - \frac{1}{2} \left\{ \tilde{P}_2^u - \tilde{P}_2^c \right\} \right] \hat{I} \otimes \Lambda_u \\ - \frac{3}{2}\tilde{P}_2^c \hat{U} \otimes \Lambda_t - \left[ \tilde{P}_1^c - \frac{\tilde{P}_2^c}{2} \right] \hat{I} \otimes \Lambda_t,$$

$$(\tilde{C}_r)_k^{ij} = \left[ \tilde{T} + \frac{3}{2}\tilde{P}_2^u - \frac{3}{2}\tilde{P}_2^c \right] \hat{U}_k^i(\Lambda_u)^j + \left[ \tilde{P}_1^u - \tilde{P}_1^c - \frac{1}{2} \left\{ \tilde{P}_2^u - \tilde{P}_2^c \right\} \right] \delta_k^i(\Lambda_u)^j \\ - \frac{3}{2}\tilde{P}_2^c \hat{U}_k^i(\Lambda_t)^j - \left[ \tilde{P}_1^c - \frac{\tilde{P}_2^c}{2} \right] \delta_k^i(\Lambda_t)^j$$

The connection between the topological decomposition and the QCD-factorization is established through

$$U_k^i(\Lambda_u)^j = \bar{H}_k^{ij}, \quad U_k^i(\Lambda_t)^j = \tilde{H}_k^{ij}, \quad (\Lambda_t)^i = \tilde{H}^i.$$


---



# QCF Factorization-Topological Equivalence

We consider the following results

$$\alpha_3^u = \alpha_3^c = \alpha_3, \quad \alpha_{3,EW}^u = \alpha_{3,EW}^c = \alpha_{3,EW}, \quad \beta_i^u = \beta_i^c = \beta_i, \quad b_i^u = b_i^c = b_i$$

$$|\alpha_{4,EW}^c - \alpha_{4,EW}^u| < 10^{-3}$$

$$|\alpha_4^c - \alpha_4^u| \sim 2\%$$

NLO

NNLO

*Bell, Beneke, Huber, Li:2002.03262*

QCDF to topological transformation rules

$$T = A_{M_1 M_2} \alpha_1,$$

$$C = A_{M_1 M_2} \alpha_2,$$

$$E = A_{M_1 M_2} \beta_1,$$

$$A = A_{M_1 M_2} \beta_2,$$

$$T_{AS} = A_{M_1 M_2} \beta_{S1},$$

$$T_{ES} = A_{M_1 M_2} \beta_{S2},$$

$$S = -A_{M_1 M_2} \left[ \alpha_3 + \beta_{S3} - \frac{\alpha_{3,EW}}{2} - \frac{\beta_{S3,EW}}{2} \right],$$

$$P = -A_{M_1 M_2} \left[ \alpha_4^c + \beta_3 - \frac{\alpha_{4,EW}^c}{2} - \frac{\beta_{3,EW}}{2} \right],$$

$$A_{M_1 M_2} = (1.25 \pm 0.17) \text{ GeV}^3$$

# Further details on the $\chi^2$ -fit

Best QCDF fit point (modulus in  $\text{GeV}^3$ )

$$A_{M_1 M_2} \alpha_1 = 1.072 + 5.596 \times 10^{-5} i,$$

$$A_{M_1 M_2} \alpha_2 = 0.136 + 0.073 i,$$

$$A_{M_1 M_2} \beta_1 = -0.117 - 0.007 i,$$

$$A_{M_1 M_2} \beta_2 = A_{M_1 M_2} \beta_1,$$

$$A_{M_1 M_2} \beta_{S1} = -0.074 - 0.0112 i,$$

$$A_{M_1 M_2} \beta_{S2} = 0.054 - 0.049 i,$$

$$A_{M_1 M_2} \alpha_{3,EW} = -0.193 - 0.045 i,$$

$$A_{M_1 M_2} \alpha_{4,EW}^c = 0.181 + 0.053 i,$$

$$A_{M_1 M_2} \beta_{3,EW} = 0.005 - 0.006 i,$$

$$A_{M_1 M_2} b_{4,EW} = A_{M_1 M_2} \beta_{3,EW},$$

$$A_{M_1 M_2} \beta_{S3,EW} = -0.188 + 0.007 i,$$

$$A_{M_1 M_2} b_{S4,EW} = 0.061 + 0.098 i,$$

$$A_{M_1 M_2} \beta_4 = -0.003 + 0.013 i,$$





$$A_{M_1 M_2} \beta_{S4} = 0.031 - 0.030 i,$$

$$A_{M_1 M_2} (\alpha_3 + \beta_{S3}) = 0.230 + 0.067 i,$$

$$A_{M_1 M_2} (\alpha_4 + \beta_3) = -0.242 - 0.062 i$$

Obtained by mapping the SU(3)-fit results into the QCDF amplitudes.

# QCF Factorization confidence regions



# Summary and Outlook

- We have established a set of transformation rules between the QCD factorization and the topological representation of physical amplitudes.
- By fitting to data we have determined bounds for different QCDF amplitudes.
- The real and imaginary components of the weak annihilation amplitudes as allowed by data can be between 4% and 30%.
- SU(3) symmetry assumed so far.
- Introduce SU(3) breaking by fitting to data the weak annihilation amplitudes combining NLO and NNLO results for independent channels

# Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 945422



# SU(3) amplitudes from data

Fit for the modulus and and phases of the relevant parameters.

Use random sampling to obtain the best fit point with  $10^9$  points:

- Calculate the  $\chi^2$  function for  $10^6$  points assuming a flat probability distribution.
- Select the best 5 points leading to the minimum  $\chi^2$ .
- Use these partial minimums as starting points for the Sequential Least Square Programming algorithm, SLSQP.
- Repeat  $10^3$  times to get the overall minimum.

To obtain the 65 % C.L regions apply a likelihood ratio test using Wilk's theorem.

# Topological decomposition

$$\begin{aligned}
 \mathcal{T}^{TDA} = & \underline{T} B_i(M)_j^i \bar{H}_k^{jl}(M)_l^k + \underline{C} B_i(M)_j^i \bar{H}_k^{lj}(M)_l^k + \underline{A} B_i \bar{H}_j^{il}(M)_k^j (M)_l^k \\
 & + \underline{E} B_i \bar{H}_j^{li}(M)_k^j (M)_l^k + \underline{T_{ES}} B_i \bar{H}_l^{ij}(M)_j^l (M)_k^k + \underline{T_{AS}} B_i \bar{H}_l^{ji}(M)_j^l (M)_k^k \\
 & + \underline{T_S} B_i(M)_j^i \bar{H}_l^{lj}(M)_k^k + \underline{T_{PA}} B_i \bar{H}_l^{li}(M)_k^j (M)_j^k + \underline{T_P} B_i(M)_j^i (M)_k^j \bar{H}_l^{lk} \\
 & + \underline{T_{SS}} B_i \bar{H}_l^{li}(M)_j^j (M)_k^k,
 \end{aligned}$$

$T$  :Color allowed tree.

$P$  : QCD-penguin.

$C$  : Color-suppressed tree.

$S$  : QCD-singlet penguin.

$E$  : W-exchange diagram.

$A$  : Annihilation.

# Further details on the $\chi^2$ -fit

## Constraints from QCDF

Taking into account  $\alpha_1(\pi\pi) = 1.000_{-0.069}^{+0.029} + (0.011_{-0.050}^{+0.023})i$ .

*Beneke Huber et al: 0911.3655*

We impose  $\Re(\alpha_1) = 1.000_{-0.138}^{+0.138}$

In addition we require

$$T_{PA} = T_{SS} = T_S = 0, \quad |T_P| < 10\%$$

## Phenomenological constraints

$$\begin{aligned} Br(B_s \rightarrow \pi^0\pi^0) &< 2.10 \times 10^{-4}, & Br(B_s \rightarrow \eta\pi^0) &< 10^{-3}, \\ Br(B^0 \rightarrow \eta\eta) &< 10^{-6}, & Br(B^0 \rightarrow \eta'\eta') &< 1.7 \times 10^{-6}, \\ Br(B^0 \rightarrow \eta'\eta) &< 1.2 \times 10^{-6}, & A_{CP}(B_s \rightarrow \eta K^0) &< 10^{-3}. \end{aligned}$$



# SU(3) amplitudes from data

Include  $\eta$  contributions in the Feldmann–Kroll–Stech scheme

$\theta_{FKS}$  mixing angle *T. Feldmann et al: 9802409*

Channel	$A_3^T$	$C_{3T}^T$	$A_6^T$	$C_6^T$	$A_{15}^T$	$C_{15}^T$	$B_3^T$	$B_6^T$	$B_{15}^T$	$D_3^T$
$B^- \rightarrow \eta_q \pi^-$	0	$\sqrt{2}$	$\sqrt{2}$	0	$3\sqrt{2}$	$2\sqrt{2}$	0	$\sqrt{2}$	$3\sqrt{2}$	$\sqrt{2}$
$B^- \rightarrow \eta_s \pi^-$	0	0	0	1	0	-1	0	1	3	1
$B^0 \rightarrow \eta_q \pi^0$	0	-1	-1	0	5	2	0	-1	5	-1
$B^0 \rightarrow \eta_s \pi^0$	0	0	0	$-\frac{1}{\sqrt{2}}$	0	$\frac{1}{\sqrt{2}}$	0	$-\frac{1}{\sqrt{2}}$	$\frac{5}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$
$B_s \rightarrow \eta_q K^0$	0	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	0	$-\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$
$B_s \rightarrow \eta_s K^0$	0	1	-1	0	-1	-2	0	-1	-1	1
$B^- \rightarrow \eta_q K^-$	0	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	$\frac{3}{\sqrt{2}}$	$\frac{5}{\sqrt{2}}$	0	$\sqrt{2}$	$3\sqrt{2}$	$\sqrt{2}$
$B^- \rightarrow \eta_s K^-$	0	1	1	0	3	-2	0	1	3	1
$B^0 \rightarrow \eta_q K^0$	0	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	0	$-\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$
$B^0 \rightarrow \eta_s K^0$	0	1	-1	0	-1	-2	0	-1	-1	1
$B_s \rightarrow \eta_q \pi^0$	0	0	-2	0	4	0	0	-2	4	0
$B_s \rightarrow \eta_s \pi^0$	0	0	0	$-\sqrt{2}$	0	$2\sqrt{2}$	0	$-\sqrt{2}$	$2\sqrt{2}$	0
$B^0 \rightarrow \eta_q \eta_q$	1	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	2	-1	1	1
$B^0 \rightarrow \eta_q \eta_s$	0	0	0	$\frac{1}{\sqrt{2}}$	0	$-\frac{1}{\sqrt{2}}$	$2\sqrt{2}$	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$
$B^0 \rightarrow \eta_s \eta_s$	1	0	1	0	-1	0	1	1	-1	0
$B_s \rightarrow \eta_q \eta_q$	1	0	0	0	1	0	2	0	2	0
$B_s \rightarrow \eta_q \eta_s$	0	0	0	0	0	$\sqrt{2}$	$2\sqrt{2}$	0	$-\sqrt{2}$	$\sqrt{2}$
$B_s \rightarrow \eta_s \eta_s$	1	1	0	0	-2	-2	1	0	-2	1