



Contribution ID : 51

Type : **not specified**

VLBI and gamma-ray studies of radio galaxies

Thursday, 5 October 2017 17:30 (15)

The γ -ray sky is strongly dominated by blazars, i.e. AGN with relativistic jets oriented closely with our line of sight. Radio galaxies are their misaligned counterparts, and make up about ~ 1 -2% of all AGN observed by Fermi-LAT. At TeV energies, only 5 radio galaxies have currently been detected, but recent work has shown that the CTA has good potential for detecting more of these elusive Very-High-Energy sources. In spite of their small numbers in γ -ray catalogs, radio galaxies provide us with a view of AGN jets which is less biased by Doppler boosting effects, and allow us to test jet production and emission models in light of the unified scheme of radio-loud AGN. The combination of γ -ray data and high-resolution Very Long Baseline Interferometry (VLBI) studies is a powerful tool in order to investigate these objects. We present selected results of an ongoing study focused on the radio galaxies in the southern-hemisphere VLBI (and multi-wavelength) monitoring program TANAMI.

Summary

Primary author(s) : Mr. ANGIONI, Roberto (Max-Planck Institut für Radioastronomie)

Co-author(s) : Dr. MUELLER, Cornelia (Radboud Uni. Nijmegen); Prof. ROS, Eduardo (MPI für Radioastronomie & Univ. de València); Prof. KADLER, Matthias (Uni. Wuerzburg); Dr. OJHA, Roopesh (NASA GSFC)

Presenter(s) : Mr. ANGIONI, Roberto (Max-Planck Institut für Radioastronomie)

Session Classification : Science symposium